
MATHEMATICS

9709/41

Paper 4

October/November 2017

MARK SCHEME

Maximum Mark: 50

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2017 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol FT implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously “correct” answers or results obtained from incorrect working.
 - Note: B2 or A2 means that the candidate can earn 2 or 0.
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking g equal to 9.8 or 9.81 instead of 10.

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF/OE	Any Equivalent Form (of answer is equally acceptable) / Or Equivalent
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
CAO	Correct Answer Only (emphasising that no “follow through” from a previous error is allowed)
CWO	Correct Working Only – often written by a ‘fortuitous’ answer
ISW	Ignore Subsequent Working
SOI	Seen or implied
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

- MR –1 A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become “follow through” marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR –2 penalty may be applied in particular cases if agreed at the coordination meeting.
- PA –1 This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Question	Answer	Marks	Guidance
1	$[12 \cos 25 = 3a]$	M1	For use of Newton's second law
	$a = 4 \cos 25 = 3.625$	A1	
	$[s = \frac{1}{2} \times 4 \cos 25 \times 5^2]$	M1	For use of $s = ut + \frac{1}{2}at^2$ OE
	Distance = 45.3 m	A1	
		4	

Question	Answer	Marks	Guidance
2(i)	Power = $1150 \times 12 = 13\,800\text{W}$	B1	For use of $P = F \times v$ Allow 13.8 kW
		1	
2(ii)	Driving force = $\frac{25000}{12}$	B1	Using $F = \frac{P}{v}$
	$\frac{25000}{12} - 1150 - 3700g \sin 4 = 3700a$	M1	For applying Newton's 2nd law up the slope, 4 terms
	$a = -0.445 \text{ m s}^{-2}$	A1	
		3	
2(iii)	$\frac{25000}{v} - 1150 - 3700g \sin 4 = 0$	M1	For stating the equation for constant v , with 3 terms, and solving for v
	$v = 6.70 \text{ m s}^{-1}$	A1	
		2	

Question	Answer	Marks	Guidance
3(i)	640×18	M1	For use of work done = $F \times d$
	Work done = 11 520 J	A1	
		2	
3(ii)	KE at start $= \frac{1}{2} \times 840 \times 14^2 = 82\,320 \text{ J}$	B1	
	PE gained = $840g \times 8 \sin 30$ $- 840g \times 10 \sin 20 = 4870 \text{ J}$	B1	
	$\frac{1}{2} \times 840 \times v^2 = 82\,320 - 11\,520 - 4870$	M1	For using work – energy equation with 4 terms and solving for v
	$v = 12.5 \text{ m s}^{-1}$	A1	
		4	

Question	Answer	Marks	Guidance
4(i)	Acceleration = $\frac{(-25)}{2.5} = -10 \text{ m s}^{-2}$	B1	AG
		1	
4(ii)	$V = -15 + 7.5 \times 4$	M1	Using $v-t$ graph OE
	$V = 15 \text{ m s}^{-1}$	A1	
		2	
4(iii)	Using $v = 0$ at $t = 4.5$ and $t = 8$	B1	
		M1	Attempting to use area to find total distance travelled
	$\frac{1}{2} \times (4.5 + 2) \times 10$ $+ \frac{1}{2} \times (8 - 4.5) \times 15$ $+ \frac{1}{2} \times (T - 8) \times 15 = 100$	M1	For setting up an equation for total distance travelled and solving for T
	$T = 13.5$	A1	
		4	

Question	Answer	Marks	Guidance
5(i)	Acceleration = 0.4 m s^{-2}	B1	
		1	
5(ii)	$\frac{100}{t^2} - 0.1t = 0$	M1	For setting $v = 0$ and solving for t
	$t = 10 \text{ s}$	A1	
		2	
5(iii)	Distance $t = 0$ to $t = 5$ is $\frac{1}{2} (1.5 + 3.5) \times 5 = 12.5$	B1	Trapezium rule or integration
	$s(t) = \int \left(\frac{100}{t^2} - 0.1t \right) dt$	M1	For integration
	$= -\frac{100}{t} - 0.05t^2 (+C)$	A1	Correct integration
	$s(10) - s(5)$	M1	Use limits 5 and 10 used or find $+C$
	Total distance = $12.5 + 6.25 = 18.75 \text{ m}$	A1	
		5	

Question	Answer	Marks	Guidance
6(i)		M1	For resolving forces (either direction)
	$X = 75 + 50 \cos 60 (= 100)$ $Y = 50 \sin 60 (= 43.3)$	A1	For both equations, unevaluated
	Resultant = $\sqrt{(100^2 + 43.3^2)} = 109 \text{ N}$	B1	
	Angle = $\arctan \left(\frac{43.3}{100} \right) = 23.4^\circ$	B1	Must state anticlockwise from the positive x-axis or show in a diagram
		4	
6(ii)	$50 \cos \alpha - F \cos 50 = 0$	B1	Resolving forces horizontally
	$50 \sin \alpha - 3F - F \sin 50 = 0$	B1	Resolving forces vertically
	$\tan \alpha = \frac{(3F + F \sin 50)}{(F \cos 50)}$	M1	For division to find θ or for using Pythagoras to find F
	$\alpha = 80.3$	A1	
	$F = 13.1$	A1	
		5	

Question	Answer	Marks	Guidance
7(i)		M1	For applying Newton's 2nd law to either particle (correct number of terms)
	$T - 0.9 g \sin 15 = 0.9a$	A1	
	$2.5 + 0.4 g \sin 25 - T = 0.4a$	A1	
	$1.3a = 1.86\dots$	M1	Solving simultaneously for a
	$a = 1.43 \text{ m s}^{-2}$	A1	
		5	

Question	Answer	Marks	Guidance
7(ii)	$F = 0.8 \times 0.4g \cos 25$	B1	
	$2.5 + 0.4 g \sin 25 - T - F = 0$	M1	For using equilibrium of forces acting on particle <i>B</i> with 4 terms
	$T - 0.9 g \sin \theta = 0$	M1	For using equilibrium of forces acting on particle <i>A</i> with 2 terms
		M1	For solving for θ
	$\theta = 8.2^\circ$	A1	
		5	